

23/55
(19)



Euröpäisches Patentamt

European Patent Office

Office européen des brevets



(11)

2002 P41
EP 0 943 687 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
22.09.1999 Bulletin 1999/38

(51) Int Cl.⁶: **C12N 15/53**, C12N 9/04,
C12P 13/06, C12R 1/13,
C12R 1/15

(21) Application number: **99100324.5**

(22) Date of filing: **12.01.1999**

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **12.01.1998 JP 375198**
11.12.1998 JP 35351398

(71) Applicant: **Ajinomoto Co., Inc.**
Tokyo 104 (JP)

(72) Inventors:
• **Suga, Mikiko, c/o Ajinomoto Co., Inc.**
Kawasaki-shi, Kanagawa-ken (JP)
• **Sugimoto, Masakazu, c/o Ajinomoto Co., Inc.**
Kawasaki-shi, Kanagawa-ken (JP)

• **Osumi, Tsuyoshi, c/o Ajinomoto Co., Inc.**
Tokyo (JP)
• **Nakamatsu, Tsuyoshi, c/o Ajinomoto Co., Inc.**
Kawasaki-shi, Kanagawa-ken (JP)
• **Hibino, Wataru, c/o Ajinomoto Co., Inc.**
Kawasaki-shi, Kanagawa-ken (JP)
• **Ito, Mika, c/o Ajinomoto Co., Inc.**
Yokkaichi-shi, Mie-ken (JP)

(74) Representative: **HOFFMANN - EITLE**
Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)

Remarks:

The applicant has subsequently filed a sequence listing and declared, that it includes no new matter.

(54) Method of producing L-serine by fermentation

(57) Disclosed is a coryneform bacterium having resistance to azaserine or β -(2-thienyl)-DL-alanine and having L-serine productivity. Also, disclosed are D-3-phosphoglycerate dehydrogenase derived from a coryneform bacterium, in which feedback inhibition by L-serine is desensitized; the D-3-phosphoglycerate dehydrogenase, which is obtainable from a coryneform bacterium having resistance to azaserine or β -(2-thienyl)-DL-alanine and having L-serine productivity; the D-3-phosphoglycerate dehydrogenase having an amino acid sequence depicted in SEQ ID NO : 12 in Sequence Listing or the sequence including substitution, addition or deletion of one or more amino acids, wherein an ami-

no acid residue corresponding to the 325th glutamic acid residue of the amino acid sequence in the SEQ ID NO : 12 is replaced with an amino acid other than glutamic acid; the D-3-phosphoglycerate dehydrogenase that has an amino acid sequence depicted in SEQ ID NO : 11 in Sequence Listing; a DNA coding for the D-3-phosphoglycerate dehydrogenase described above; the DNA that has a base sequence depicted in SEQ ID NO : 13 in Sequence Listing; a coryneform bacterium which harbors a recombinant DNA containing the DNA; and a method of producing L-serine, comprising the steps of cultivating the bacterium described above in a medium to allow accumulation of L-serine in the medium, and collecting the L-serine from the medium.

Description

FIELD OF THE INVENTION

[0001] The present invention relates to a method of producing L-serine for use in the production of amino acid mixtures utilized in the field of pharmaceuticals, chemicals, and cosmetics, to coryneform bacteria constituting the method, to D-3-phosphoglycerate dehydrogenase (hereafter, sometimes referred to as "3-PGDH"), and to DNA coding for the 3-PGDH.

BACKGROUND OF THE INVENTION

[0002] As a conventional method of producing L-serine by fermentation, there has been reported the method in which a bacterial strain capable of converting glycine and sugar into L-serine is used in a medium containing 30 g/L of glycine to produce at most 14 g/L of L-serine. The conversion yield of glycine into L-serine by this method amounted to 46% (Kubota K. Agricultural Biological Chemistry, 49, 7-12 (1985)). Using a bacterial strain capable of converting glycine and methanol into L-serine, 53 g/L of L-serine can be produced from 100 g/L of glycine (T. Yoshida et al., Journal of Fermentation and Bioengineering, Vol. 79, No. 2, 181-183, 1995). In the method using a bacterium belonging to the genus *Nocardia*, it has been known that the L-serine productivity of the bacterium can be improved by breeding those strains resistant to serine hydroxamate, azaserine or the like (Japanese Patent Publication No. 57-1235). However, these methods involve use of glycine that is a precursor of L-serine and include complicated operation and is disadvantageous from the viewpoint of costs.

[0003] As strains that can ferment L-serine directly from a sugar and do not need addition of the precursor of L-serine to the medium, there has been known *Corynebacterium glutamicum* that is resistant to D-serine, α -methylserine, o-methylserine, isoserine, serine hydroxamate, and 3-chloroalanine but the accumulation of L-serine is as low as 0.8 g/L (Nogei Kagaku Kaishi, Vol. 48, No. 3, p201-208, 1974). Accordingly, a further strain improvements of are needed for direct fermentation of L-serine on an industrial scale.

[0004] On the other hand, regarding coryneform bacteria, there have been disclosed a vector plasmid that is capable of autonomous replication in the cell and having a drug resistance marker gene (cf. U. S. Patent 4,514,502) and a method of introducing a gene into the cell (Japanese Patent Application Laid-open No. 2-207791), and the possibility of growing L-threonine or L-isoleucine producing bacteria (U. S. Patents 4,452,890 and 4,442,208). Also, regarding the growth of L-lysine producing bacteria, there has been known a technology involving the incorporation of a gene participating in the biosynthesis of L-lysine into a vector plasmid and the amplification of the plasmid in the cell (Japanese Patent Application Laid-open No. 56-160997).

[0005] In the case of *Escherichia coli*, the enzymes participating in the biosynthesis of L-serine include an enzyme that is susceptible to feedback inhibition relative to L-serine production in the wild type and an example has been known in which the introduction of a mutant gene that has been mutated so that the feedback inhibition could be desensitized resulted in an enhancement in the L-serine (Japanese Patent No. 2584409). As such genes, there has been known specifically 3-PGDH gene (hereafter, the gene coding for 3-PGDH protein will also be referred to "serA").

[0006] Further, in the case of coryneform bacteria, an example has been known in which the amplification of 3-PGDH gene influences the productivity of L-tryptophane (Japanese Patent Application Laid-open No. 3-7591).

SUMMARY OF THE INVENTION

[0007] An object of the present invention is to provide a microorganism that converts a sugar into L-serine and to provide a method of accumulating L-serine in a culture medium utilizing the ability of the microorganism to convert the sugar into L-serine, i.e., a method of producing L-serine that is advantageous in practicing on an industrial scale.

[0008] As a result of intensive investigation on the method of producing L-serine with view to achieving the above object, it has now been discovered by the present inventors that screening a coryneform bacterium having L-serine productivity, particularly preferably a mutant strain exhibiting resistance to azaserine or β -(2-thienyl)-DL-alanine derived from a strain of the coryneform bacterium but is deficient in L-serine decomposing activity as a parent strain and conducting L-serine fermentation using the screened strain will enhance the accumulation of L-serine drastically. The present invention has been completed based on this discovery.

[0009] That is, the present invention relates to a coryneform bacterium having resistance to azaserine or β -(2-thienyl)-DL-alanine and having L-serine productivity.

[0010] Further, the present invention relates to D-3-phosphoglycerate dehydrogenase derived from a coryneform bacterium, in which feedback inhibition by L-serine is desensitized; to the D-3-phosphoglycerate dehydrogenase as described above, obtainable from a coryneform bacterium having resistance to azaserine or β -(2-thienyl)-DL-alanine and having L-serine productivity; to D-3-phosphoglycerate dehydrogenase having an amino acid sequence amino acid

sequence depicted in SEQ ID NO : 12 in Sequence Listing or the sequence including substitution, addition or deletion of one or more amino acids, wherein an amino acid residue corresponding to the 325th glutamic acid residue of the amino acid sequence in the SEQ ID NO : 12 is replaced with an amino acid other than glutamic acid; and to the D-3-phosphoglycerate dehydrogenase as described above that has an depicted in SEQ ID NO : 11 in Sequence Listing.

[0011] Still further, the present invention relates to a DNA coding for the D-3-phosphoglycerate dehydrogenase described above and to the DNA described above having a base sequence as depicted in SEQ ID NO : 13 in Sequence Listing.

[0012] Yet further, the present invention relates to a coryneform bacterium that harbors a recombinant DNA containing the DNA described above.

[0013] Further, the present invention relates to a method of producing L-serine, comprising the steps of cultivating the bacterium as described above in a medium to allow accumulation of L-serine in the medium, and collecting the L-serine from the medium.

[0014] Specific examples of the coryneform bacterium having resistance to azaserine or β -(2-thienyl)-DL-alanine and having L-serine productivity include Brevibacterium flavum AJ13324 and AJ13327 or Brevibacterium flavum AJ13325.

[0015] The present invention provides coryneform bacteria that produce L-serine from a sugar. The coryneform bacteria can be utilized in a method of producing L-serine that is industrially advantageous.

BRIEF DESCRIPTION OF THE DRAWINGS

[0016] Fig. 1 illustrates a manner of feedback inhibition of 3-PGDH derived from various strains by L-serine. The horizontal axis indicates the concentration of L-serine in the enzyme solution. The vertical axis indicates percentage of the 3-PGDH activity in the presence of L-serine to that in the absence of L-serine. Symbol \diamond illustrates a manner of feedback inhibition of 3-PGDH derived from ATCC14067 strain by L-serine. Symbol \blacksquare illustrates a manner of feedback inhibition of 3-PGDH derived from AJ13377 strain by L-serine. Symbol \blacktriangle illustrates a manner of feedback inhibition of 3-PGDH derived from AJ13324 strain by L-serine. Symbol \times illustrates a manner of feedback inhibition of 3-PGDH derived from AJ13325 strain by L-serine. Symbol $*$ illustrates a manner of feedback inhibition of 3-PGDH derived from AJ13327 strain by L-serine.

[0017] Fig. 2 illustrates the construction of plasmids pVK7 and pVK6.

DETAILED DESCRIPTION OF THE INVENTION

[0018] The coryneform bacteria referred to in the present invention are a group of microorganisms as defined in Bergey's Manual of Determinative Bacteriology, 8th ed., p. 599 (1974), which are aerobic Gram-positive rods having no acid resistance and no spore-forming ability. The coryneform bacteria include bacteria belonging to the genus Corynebacterium, bacteria belonging to the genus Brevibacterium having been hitherto classified into the genus Brevibacterium but united as bacteria belonging to the genus Corynebacterium at present, and bacteria belonging to the genus Brevibacterium closely relative to bacteria belonging to the genus Corynebacterium and bacteria belonging to the genus Microbacterium.

[0019] The coryneform bacteria of the present invention that have resistance to azaserine or β -(2-thienyl)-DL-alanine and having L-serine productivity, preferably those strains of these coryneform bacteria that are deficient in L-serine decomposing activity, are artificially mutated or induced from wild type or L-serine producing coryneform bacteria as a parent strain.

[0020] The coryneform bacteria that have resistance to azaserine or β -(2-thienyl)-DL-alanine, which are deficient in L-serine decomposing activity and have L-serine productivity can be obtained, for example, as follows. That is, Brevibacterium flavum ATCC 14067 is subjected to mutation treatment by a conventional method (contact with N-methyl-N'-nitro-N-nitroso-guanidine or the like) to obtain a mutant strain deficient in L-serine decomposing activity and using this strain as a parent strain, strains resistant to azaserine or β -(2-thienyl)-DL-alanine are obtained. From among the mutant strains obtained by the method as described above can be obtained strains that accumulate L-serine in high concentrations.

[0021] The strains resistant to azaserine or β -(2-thienyl)-DL-alanine can also be obtained by introducing the mutant serA described below into parent strains or mutant strains deficient in L-serine decomposing activity.

[0022] The term "azaserine resistance" refers to the property that a bacterium grows faster than the wild type in a medium containing azaserine. For example, those strains that form colonies on a solid medium containing 0.25 g/L of azaserine at 30°C within 4 to 5 days are said to have azaserine resistance.

[0023] Similarly, the term " β -(2-thienyl)-DL-alanine resistance" refers to the property that a bacterium grows faster than the wild type in a medium containing β -(2-thienyl)-DL-alanine. For example, those strains that form colonies on a solid medium containing 0.25 g/L of β -(2-thienyl)-DL-alanine at 30°C within 4 to 5 days are said to have β -(2-thienyl)-

DL-alanine resistance.

[0024] 3-PGDH catalyzes reaction in which 3-phosphoglycerate is oxidized into 3-phosphohydroxypyruvic acid in the presence of nicotinamide adenine dinucleotide (NAD) as a coenzyme.

[0025] The activity of 3-PGDH can be determined, for example, by the measurement of a decrease in coenzyme NADH₂ in a reverse reaction (E. Sugimoto and L. I. Pizer, JBC, 243, 2081, 1968) or synthesis of coenzyme NADH₂ in a forward reaction (Salach H. J. Method in Enzymology, vol.9, 216-220, 1966) by absorbance at 340 nm.

[0026] 3-PGDH can be purified by collecting cells from the culture broth of a coryneform bacterium, fragmenting the collected cells by sonication and subsequent ultracentrifugation, and isolating the targeted enzyme from the supernatant by a conventional method. More particularly, 3-PGDH can be purified by sequentially concentrating fractions having 3-PGDH activity by precipitation with ammonium sulfate, gel filtration, cation exchange resin chromatography, anion exchange resin chromatography, reverse phase chromatography and the like.

[0027] 3-PGDH derived from a wild type coryneform bacterium is susceptible to feedback inhibition by L-serine and its activity is almost completely inhibited in the presence of 10 mM of L-serine. By the term "3-PGDH in which feedback inhibition by L-serine is desensitized" is meant 3-PGDH having 20% or more, preferably 40% or more, more preferably 90% or more of the activity in the absence of L-serine even in the presence of 10 mM of L-serine. 3-PGDH derived from Brevibacterium flavum AJ13327 described in the examples hereinbelow retains substantially 100% of the activity in the presence of 80 mM of L-serine and therefore one of the most preferred 3-PGDHs.

[0028] 3-PGDH derived from a wild type coryneform bacterium (hereafter, DNA coding for this is also referred to as "wild type serA") has the amino acid sequence depicted in SEQ ID NO : 12 in Sequence Listing. Specific examples of the 3-PGDH in which feedback inhibition by L-serine is desensitized (hereafter, DNA coding for this is also referred to as "mutant serA") include D-3-phosphoglycerate dehydrogenase characterized in that in D-3-phosphoglycerate dehydrogenase having the amino acid sequence depicted in SEQ ID NO : 12 in Sequence Listing or the same amino acid sequence as above but has substitution, addition or deletion of one or more amino acids, the amino acid residue corresponding to the 325th glutamic acid residue of the amino acid sequence in the SEQ ID NO : 12 has been substituted by other amino acid. Most preferred as the other amino acid residue is a lysine residue.

[0029] The DNA fragment containing serA gene from a coryneform bacterium can be isolated, for example, by preparing chromosomal DNA according to the method of Saito and Miura (H. Saito and K. Miura, Biochem. Biophys. Acta, 72, 619 (1963)) or the like and then amplifying serA gene by polymerase chain reaction method (PCR: polymerase chain reaction; cf. White, T. J. et al.; Trends Genet. 5, 185 (1989)). For example, in order to amplify DNA fragment containing ORF (172 to 1705) of SEQ ID NO : 11 in Sequence Listing, any 20 to 30 bases are selected from the region from the first base in SEQ ID NO : 11 to the base immediately before ATG to obtain one primer. Further, any 20 to 30 bases are selected from the region from the base immediately after the termination codon to the last base in SEQ ID NO : 11 to obtain another primer.

[0030] When serA is isolated from a wild type strain of 3-PGDH, wild type serA is obtained and isolation of serA from a mutant harboring 3-PGDH in which feedback inhibition by L-serine is desensitized (3-PGDH mutant) gives mutant serA. Specifically, the wild type serA has the sequence depicted in SEQ ID NO : 11 in Sequence Listing, and mutant serA has the sequence depicted in SEQ ID NO : 13 in Sequence Listing.

[0031] It is preferred that serA amplified by PCR method is ligated with vector DNA autonomously replicable in cells of Escherichia coli and/or coryneform bacteria to prepare recombinant DNA, and the recombinant DNA is introduced into cells of Escherichia coli beforehand. Such provision makes following operations easy. The vector autonomously replicable in cells of Escherichia coli is preferably a plasmid vector which is preferably autonomously replicable in cells of a host, including, for example, pUC19, pUC18, pBR322, pHSG299, pHSG399, pHSG398, and RSF1010.

[0032] Recombinant DNA may be prepared by utilizing transposon (WO 02/02627 International Publication Pamphlet, WO 93/18151 International Publication Pamphlet, European Patent Application Laid-open No. 0445385, Japanese Patent Application Laid-open No. 6-46867, Vertes, A. A. et al., Mol. Microbiol., 11, 739-746 (1994), Bonamy, C., et al., Mol. Microbiol., 14, 571-581 (1994), Vertes, A. A. et al., Mol. Gen. Genet., 245, 397-405 (1994), Jagar, W. et al., FEMS Microbiology Letters, 126, 1-6 (1995), Japanese Patent Application Laid-open No. 7-107976, Japanese Patent Application Laid-open No. 7-327680, etc.), phage vectors, recombination of chromosomes (Experiments in Molecular Genetics, Cold Spring Harbor Laboratory Press (1972); Matsuyama, S. and Mizushima, S., J. Bacteriol., 162, 1196 (1985)) and the like.

[0033] When a DNA fragment having an ability to allow a plasmid to be autonomously replicable in coryneform bacteria is inserted into these vectors, they can be used as a so-called shuttle vector autonomously replicable in both Escherichia coli and coryneform bacteria.

[0034] Such a shuttle vector includes the followings. Microorganisms harboring each of vectors and deposition numbers in international deposition facilities are shown in parentheses. pH4: Escherichia coli AJ12617 (FERM BP-3532) pAJ655: Escherichia coli AJ11882 (FERM BP-136)

Corynebacterium glutamicum SR8201 (ATCC 39135)

pAJ1844: Escherichia coli AJ11883 (FERM BP-137)

Corynebacterium glutamicum SR8202 (ATCC 39136)

pAJ611: Escherichia coli AJ11884 (FERM BP-138)

pAJ3148: Corynebacterium glutamicum SR8203 (ATCC 39137)

pAJ440: Bacillus subtilis AJ11901 (FERM BP-140)

[0035] These vectors are obtainable from the deposited microorganisms as follows. Cells collected at a logarithmic growth phase were lysed by using lysozyme and SDS, followed by separation from a lysate by centrifugation at 30,000 x g to obtain a supernatant to which polyethylene glycol is added, followed by fractionation and purification by means of cesium chloride-ethidium bromide equilibrium density gradient centrifugation.

[0036] Escherichia coli can be transformed by introducing a plasmid in accordance with, for example, a method of D. M. Morrison (Methods in Enzymology, 68, 326 (1979)) or a method in which recipient cells are treated with calcium chloride to increase permeability for DNA (Mandel, M. and Higa, A., J. Mol. Biol., 53, 159 (1970)).

[0037] Introduction of plasmids to coryneform bacteria to cause transformation can be performed by the electric pulse method (Sugimoto et al., Japanese Patent Application Laid-open No. 2-207791).

[0038] Examples of the coryneform bacterium used to introduce the DNA described above include, for example, the following wild type strains:

Corynebacterium acetoacidophilum ATCC 13870;

Corynebacterium acetoglutamicum ATCC 15806;

Corynebacterium callunae ATCC 15991;

Corynebacterium glutamicum ATCC 13032;

(Brevibacterium divaricatum) ATCC 14020;

(Brevibacterium lactofermentum) ATCC 13869;

(Corynebacterium lilium) ATCC 15990;

(Brevibacterium flavum) ATCC 14067;

Corynebacterium mellesecola ATCC 17965;

Brevibacterium saccharolyticum ATCC 14066;

Brevibacterium immariophilum ATCC 14068;

Brevibacterium roseum ATCC 13825;

Brevibacterium thiogenitalis ATCC 19240;

Microbacterium ammoniaphilum ATCC 15354;

Corynebacterium thermoaminogenes AJ12340 (FERM BP-1539).

[0039] The transformed strains obtained by introduction of recombinant DNA containing the mutant serA into the coryneform bacterial as described above produces 3-PGDH in which feedback inhibition by L-serine is desensitized. The transformed strains have resistance to azaserine or β -(2-thienyl)-DL-alanine.

[0040] For L-serine production using the strain of the present invention, the following methods may be used. As the medium to be used, there can be used conventional liquid mediums containing carbon sources, nitrogen sources, inorganic salts, and optionally organic trace nutrients such as amino acids, vitamins, etc., if desired.

[0041] As carbon sources, it is possible to use sugars such as glucose, sucrose, fructose, galactose; saccharified starch solutions, sweet potato molasses, sugar beet molasses and highest molasses which are including the sugars described above; organic acids such as acetic acid; alcohols such as ethanol; glycerol and the like.

[0042] As nitrogen sources, it is possible to use ammonia gas, aqueous ammonia, ammonium salts, urea, nitrates and the like. Further, organic nitrogen sources for supplemental use, for example, oil cakes, soybean hydrolysate liquids, decomposed casein, other amino acids, corn steep liquor, yeast or yeast extract, peptides such as peptone, and the like, may be used.

[0043] As inorganic ions, phosphoric ion, magnesium ion, calcium ion, iron ion, manganese ion and the like may be added optionally.

[0044] In case of using the microorganism of the present invention which requires nutrients such as amino acids for its growth, the required nutrients should be supplemented.

[0045] The microorganisms are incubated usually under aerobic conditions at pH 5 to 8 and temperature ranges of 25 to 40°C. The pH of the culture medium is controlled at a predetermined value within the above-described ranges depending on the presence or absence of inorganic or organic acids, alkaline substances, urea, calcium carbonate, ammonia gas, and the like.

[0046] L-Serine can be collected from the fermentation liquid, for example, by separating and removing the cells, subjecting to ion exchange resin treatment, concentration cooling crystallization, membrane separation, and other known methods in any suitable combination. In order to remove impurities, activated carbon adsorption and recrystallization may be used for purification.

DESCRIPTION OF PREFERRED EMBODIMENTS

(Example 1) Construction of novel L-serine producing bacteria Brevibacterium flavum AJ13324 and AJ13327

[0047] Brevibacterium flavum AJ13324 and AJ13327 were constructed from Brevibacterium flavum AJ13377 that is deficient in L-serine decomposing activity obtained from wild type strain Brevibacterium flavum ATCC 14067.

[0048] To obtain a mutant, cells proliferated for 24 hours in a bouillon medium (a medium containing 1 g of fish meat extract, 1 g of polypeptone, 0.5 g of yeast extract, and 0.5 g of sodium chloride in 1 liter of water, adjusted to pH 7.0) were suspended in 100 mM phosphate buffer (pH 7.0) (containing 10^9 to 10^{10} cells/ml). NG (N-methyl-N'-nitro-N-nitrosoguanidine) was added to the suspension to a concentration of 200 µg/ml and left to stand at 30°C for 30 minutes. The thus NG treated cells were washed well with the above-described buffer.

[0049] To select strains having no L-serine decomposing activity from the NG treated cells, NG treated cells of Brevibacterium flavum ATCC 14067 after washed were spread on a bouillon agar medium and incubated at 30°C for 24 hours to allow colony formation. Then, the colonies on the bouillon agar medium were used as a negative and replica formation was performed on a minimal medium and a minimal medium for selection. Then, strains were screened that grow on the minimal medium but do not grow on the minimal medium for selection. The minimal medium was a medium that contained 20 g of glucose, 1 g of ammonium sulfate, 1 g of potassium dihydrogen phosphate, 2.5 g of urea, 0.4 g of magnesium sulfate heptahydrate, 0.01 g of iron (II) sulfate heptahydrate, 0.01 g of manganese sulfate tetra- to pentahydrate, 50 µg of biotin, 200 µg of thiamin hydrochloride, 200 µg of nicotinic acid amide, and 2.0 g of agar per liter of distilled water. The minimal medium for selection was a medium that contained 1 g of ammonium sulfate, 1 g of potassium dihydrogen phosphate, 2.5 g of urea, 0.4 g of magnesium sulfate heptahydrate, 0.01 g of iron (II) sulfate heptahydrate, 0.01 g of manganese sulfate tetra- to pentahydrate, 50 µg of biotin, 200 µg of thiamin hydrochloride, 200 µg of nicotinic acid amide, 0.5 g of L-serine and 2.0 g of agar per liter of distilled water. Among the mutants obtained by this method were found many strains that have no L-serine decomposing activity and Brevibacterium flavum AJ13377 was obtained as one of such strains.

[0050] To select azaserine resistant strains from NG treated strains using Brevibacterium flavum AJ13377 as a parent strain, NG treated Brevibacterium flavum AJ13377 cells after washed were inoculated on a minimal medium for selection. The minimal medium for selection was a medium that contained 20 g of glucose, 1 g of ammonium sulfate, 1 g of potassium dihydrogen phosphate, 2.5 g of urea, 0.4 g of magnesium sulfate heptahydrate, 0.01 g of iron (II) sulfate heptahydrate, 0.01 g of manganese sulfate tetra- to pentahydrate, 50 µg of biotin, 200 µg of thiamin hydrochloride, 200 µg of nicotinic acid amide, and 250 mg of azaserine per liter of distilled water. The NG treated mutant was incubated in the above-described medium at 30°C for 5 to 10 days. The cell culture thus obtained was spread on a bouillon agar medium and incubated at 30°C for 24 hours for colony formation. Azaserine resistant strains were obtained from the strains that formed colonies. The mutants thus obtained included many strains that accumulated L-serine in considerable amounts at high yields. From the strains were obtained two strains, i.e., Brevibacterium flavum AJ13324 and AJ13327. It was confirmed that these strains were able to grow in the presence of 0.25 g/L of azaserine.

(Example 2) Construction of novel L-serine producing bacterium Brevibacterium flavum AJ13325

[0051] Brevibacterium flavum AJ13325 was constructed from Brevibacterium flavum AJ13377 lacking L-serine decomposing activity, which was obtained from the wild type strain Brevibacterium flavum ATCC 14067.

[0052] To select β-(2-thienyl)-DL-alanine resistant strains from NG treated strains using Brevibacterium flavum AJ13377 as a parent strain, Brevibacterium flavum AJ13377 cells were NG treated and washed before their inoculation on a minimal medium for selection. The minimal medium for selection was a medium that contained 20 g of glucose, 1 g of ammonium sulfate, 1 g of potassium dihydrogen phosphate, 2.5 g of urea, 0.4 g of magnesium sulfate heptahydrate, 0.01 g of iron (II) sulfate heptahydrate, 0.01 g of manganese sulfate tetra- to pentahydrate, 50 µg of biotin, 200 µg of thiamin hydrochloride, 200 µg of nicotinic acid amide, and 250 mg of β-(2-thienyl)-DL-alanine per liter of distilled water. The NG treated mutant was incubated in the above-described medium at 30°C for 5 to 10 days. The cell culture thus obtained was spread on a bouillon agar medium and incubated at 30°C for 24 hours for colony formation. β-(2-Thienyl)-DL-alanine resistant strains were obtained from the strains that formed colonies. The mutants thus obtained included many strains that accumulated L-serine in considerable amounts at high yields. Brevibacterium flavum AJ13325 was obtained as one of such strains. It was confirmed that these strains were able to grow in the presence of 0.25 g/L of β-(2-thienyl)-DL-alanine.

(Example 3) Production of L-serine by novel L-serine producing bacteria Brevibacterium flavum AJ13324, AJ13325 and AJ13327

[0053] Brevibacterium flavum AJ13324, AJ13325 and AJ13327 were each incubated on a bouillon agar medium at

30°C for 24 hours and a loopful of each microorganism was inoculated in a 500 ml shaking flask containing 20 ml of a fermentation medium having the composition shown in Table 1. As a control, the parent strains Brevibacterium flavum ATCC 14067 and AJ13377 were inoculated in the same manner as described above. The medium was adjusted to pH 7.0 with potassium hydroxide and autoclaved at 115°C for 15 minutes. After the sterilization and cooling, calcium carbonate that had been dry air sterilized at 180°C for 3 hours was added in an amount of 5 g/L.

Table 1

Component	Content/liter
Glucose	110.0 g
Potassium dihydrogen phosphate	0.4 g
Magnesium sulfate heptahydrate	0.4 g
Iron (II) sulfate heptahydrate	0.01 g
Manganese sulfate tetra- to pentahydrate	0.01 g
Ammonium sulfate	25.0 g
Thiamin hydrochloride	100 µg
Biotin	100 µg
Soy bean protein hydrochloric acid hydrolysate ("Mieki" (registered trademark))	40 ml
pH	7.0

[0054] Determination of L-serine using high performance liquid chromatography (Hitachi L-8500 Amino Acid Auto-analyzer) revealed that Brevibacterium flavum AJ13324, AJ13325 and AJ13327 accumulated L-serine in the medium in amounts of 15.2 g/L, 14.3 g/L, and 15.4 g/L, respectively. On the other hand, Brevibacterium flavum strains ATCC 14067 and AJ13377 incubated as a control accumulated L-serine in amounts of 0 g/L and 5.0 g/L, respectively.

[0055] The culture broth of Brevibacterium flavum AJ13324 was centrifuged and the supernatant was subjected to desalting treatment using cation exchange resin, followed by chromatographic separation with cation exchange resin and anion exchange resin to remove byproducts and purification by crystallization to obtain L-serine crystals of at least 99% purity at a yield from broth of 55%.

(Example 4) Measurement of 3-PGDH activity

[0056] Brevibacterium flavum AJ13324, AJ13325 and AJ13327 were each incubated on a bouillon agar medium at 30°C for 24 hours and a loopful of each microorganism was inoculated in a 500 ml shaking flask containing 50 ml of a fermentation medium having the composition shown in Table 2. As a control, the parent strains Brevibacterium flavum ATCC 14067 and AJ13377 were inoculated in the same manner as described above. The medium for inoculation was adjusted to pH 5.5 with sodium hydroxide and autoclaved at 115°C for 15 minutes.

Table 2

Component	Content/liter
Glucose	30.0 g
Potassium dihydrogen phosphate	1.0 g
Magnesium sulfate heptahydrate	0.4 g
Iron (II) sulfate heptahydrate	0.01 g
Manganese sulfate tetra- to pentahydrate	0.01 g
Ammonium sulfate	3.0 g
Soy bean protein hydrochloric acid hydrolysate ("Mieki" (registered trademark))	3.0 ml
Thiamin hydrochloride	200 µg
Biotin	50 µg
Urea	3.0 g
Yeast extract	2.0 g
pH	5.5

[0057] After collecting cells from the culture broth of each strain, the cells were washed twice with physiological saline and suspended in 50 mM sodium phosphate buffer (pH 7.0) containing 2 mM dithiothreitol. After ice cooling, the sus-

pension was subjected to a sonicator to fragment the cells and the resulting liquid was ultracentrifuged. The ultracentrifugation was run at 45,000 rpm for 1 hour to obtain a crude enzyme solution.

[0058] The enzyme activity of 3-PGDH was measured by the method of Salach H. J. et al. (Method in Enzymology, vol 9. 216-220 (1966)).

[0059] More specifically, 0.4 ml of 0.015 M NAD, 0.12 ml of 0.25 M EDTA (pH 9, NaOH), 0.1 ml of 0.05 M glutathione (pH 6, KOH), 0.5 ml of 1 M hydrazine (pH 9, acetate), 0.6 ml of 1 M Tris (pH 9, HCl), a suitable concentration of L-serine (0 to 40 mM), and water to make 2.3 ml, warmed to 25°C in advance, were added. Then, 0.2 ml of the crude enzyme solution was added and the temperature was kept the same for 5 minutes. Thereafter, 0.5 ml of 0.1 M 3-PGA (3-phosphoglycerate disodium salt, pH 7, NaOH) was added. After stirring, the absorbance at 340 nm of the reaction mixture was measured for 30 seconds. The reaction was carried out at 25°C.

[0060] For the measurement of activity, Hitachi U-2000A spectrophotometer was used.

[0061] Fig. 1 illustrates the results obtained. AJ13377 strain was relieved of L-serine sensitivity as compared with the wild type strain ATCC 14067. The AJ13324 strain was more relieved of L-serine sensitivity and the AJ13325 strain was of the same level as the AJ13324 strain in this respect. The AJ13327 strain was relieved of L-serine sensitivity greatly. And the inhibition was completely desensitized even in the presence of 80 mM L-serine.

[0062] Although some examples of desensitization of the inhibition of 3-PGDH by L-serine were reported on Escherichia coli (Tosa and Pizer, J. Bacteriol. 106: 972-982 (1971) or Japanese Patent Application Laid-open No. 6-510911), there has been known no example of complete desensitization of the inhibition in the presence of such a high concentration of L-serine.

(Example 5) Cloning of coryneform bacteria-derived wild type and mutant serA

[0063] As shown in Example 4, the feedback inhibition by L-serine was completely desensitized in the AJ13327 strain. Accordingly, cloning of serA gene coding for wild type 3-PGDH derived from the ATCC 14067 strain and mutant 3-PGDH derived from the AJ13327 strain was attempted in order to elucidate what the variation was like and confirm the amplification effect of 3-PGDH.

[0064] To amplify serA from the chromosome of Brevibacterium flavum using a PCR method, it is necessary to make a corresponding primer. Since no report has been made on the cloning and nucleotide sequence of serA of Brevibacterium flavum, the sequence of serA derived from Corynebacterium was used. Plasmid pDTS9901 was extracted from the strain Corynebacterium glutamicum K82 (cf. FERM BP-2444 and Japanese Patent Application Laid-open No. 3-7591) in which the serA fragment derived from Corynebacterium was cloned using Wizard Minipreps DNA Purification System (manufactured by Promega) and a DNA fragment of about 1.4 kb containing serA was cleaved with restriction enzyme BamHI (manufactured by Takara Shuzo Co., Ltd.).

[0065] As a vector for cloning the gene fragment, there was used a newly constructed cloning vector pVK7 for coryneform bacteria.

[0066] pVK7 was constructed by ligating (a cloning vector for Escherichia coli) pHSG299 (Kmr; Takeshita, S. et al., Gene, 61, 63-74 (1987), Japanese Patent Application Laid-open No. 10-215883), to pAM330, a cryptic plasmid of Brevibacterium lactofermentum, in the manner described below. pHSG299 was cleaved with monospecific restriction enzyme AvaII (manufactured by Takara Shuzo Co., Ltd.) and blunt ended with T4 DNA polymerase. This was ligated with pAM330 that had been cleaved with HindIII (manufactured by Takara Shuzo Co., Ltd.) and blunt ended with T4 DNA polymerase. The two types of plasmids obtained were designated pVK6 and pVK7 depending on the direction of pAM330 insertion relative to pHSG299, and pVK7 was used in the following experiments. pVK7 was capable of autonomous replication in Escherichia coli and Brevibacterium lactofermentum and retains the multiple cloning site and lacZ' derived from pHSG299. Fig. 2 illustrates the process of constructing pVK6 and pVK7.

[0067] To the shuttle vector pVK7 thus constructed was ligated a DNA fragment of about 1.4 kb containing serA. pDTS9901 was cleaved with restriction enzyme BamHI (manufactured by Takara Shuzo Co., Ltd.) and ligated to pVK7 also cleaved with restriction enzyme BamHI. The ligation of DNA was performed using DNA Ligation Kit (manufactured by Takara Shuzo Co., Ltd.) according to the prescribed method.

[0068] For the sequencing reaction, use was made of PCR thermal cycler MP type (manufactured by Takara Shuzo Co., Ltd.) and of Dye Terminator Cycle Sequencing FS Ready Reaction Kit (manufactured by Perkin Elmer). As the DNA primer, there were used M13(-21), RV primer (manufactured by Takara Shuzo Co., Ltd.). The SEQ ID NO: 1 in Sequence Listing shows the sequence thus obtained. SEQ ID NO: 2 shows an amino acid sequence that can be coded for by this sequence.

[0069] A primer was synthesized based on the base sequence thus determined and serA was amplified by a PCR method using the chromosomal DNA of the mutant Brevibacterium flavum AJ13327 as a template. The SEQ ID NOS: 3 and 4 in Sequence Listing show the N-terminal side and C terminal side sequences, respectively, of the DNA primer that were synthesized for gene amplification.

[0070] In the preparation of the chromosomal DNA of Brevibacterium flavum, use is made of Genomic DNA Purifi-

cation Kit (Bacterial) (manufactured by Advanced Genetic Technologies Corp.) and the preparation method was according to the annexed protocol.

[0071] For the PCR reaction, use is made of PCR Thermal Cycler MP type (Takara Shuzo Co., Ltd.) and of TaKaRa Taq (manufactured by Takara Shuzo Co., Ltd.).

[0072] The PCR product was ligated directly to plasmid pCR2.1 vector using Original TA Cloning Kit (manufactured by Invitrogen) and transformation was performed using competent cell of INV α F'. The transformed cells were spread on L medium (10 g/L of bactotryptone, 5 g/L of bacto-yeast extract, 15 g/L of NaCl, and 15 g/L of agar) further containing 40 μ g/ml of X-Gal (5-bromo-4-chloro-3-indolyl- β -D-galactoside) and 25 μ g/ml of Kanamycin, and incubated overnight. The white colonies, which appeared, were collected and separated to single colonies to obtain a transformed strain.

[0073] Plasmids were extracted from the transformed strain and those plasmids of which insertion of the serA fragment was confirmed by a PCR method were treated with restriction enzyme EcoRI and ligated to the shuttle vector pVK. Determination of the base sequence of the product suggested that no full-length sequence be contained on the C-terminal side. The sequence thus obtained corresponds to the region from 277 bases upstream of SEQ ID NO : 13 on the 5' side to the 1134th base of SEQ ID NO : 13 in Sequence Listing on the 3' side.

[0074] To obtain a fragment containing the full length serA gene, cloning of a deleted part from the chromosomal DNA of Brevibacterium flavum AJ13327 strain was performed according to the annexed protocol using TaKaRa LA PCR in vitro Cloning Kit (manufactured by Takara Shuzo Co., Ltd.)

[0075] First, the chromosomal DNA thus prepared was completely digested with various restriction enzymes and ligated with cassettes having respective restriction enzyme sites corresponding thereto. Cassette primer (C1) (SEQ ID NO : 5 in Sequence Listing) and a primer complementary to a known region of DNA (S1) (SEQ ID NO : 6 in Sequence Listing) were used for carrying out first PCR. Using a portion of the reaction mixture, second PCR was carried out with inner primer C2 (SEQ ID NO : 7 in Sequence Listing) and S2 (SEQ ID NO : 8 in Sequence Listing) to amplify only the targeted DNA.

[0076] When EcoRI (manufactured by Takara Shuzo Co., Ltd.) was used as the restriction enzyme, the amplification of the targeted DNA was confirmed and the base sequence of the PCR product was determined directly. Based on the base sequence thus obtained, a primer coding for the C-terminal side was made and the fragments containing full length serA were collected from Brevibacterium flavum ATCC 14067 as a wild type strain and Brevibacterium flavum AJ13327 as a mutant strain. SEQ ID NOS : 9 and 10 in Sequence Listing show the sequences of N-terminal and C-terminal side DNA primers, respectively.

[0077] The gene fragments containing wild type serA and mutant serA, respectively, in their full length were ligated to EcoRI-cleaved shuttle vector pVK7 using Original TA Cloning Kit (manufactured by Invitrogen). Plasmids harboring respective gene fragments were made separately and their base sequence was determined. SEQ ID NOS : 11 and 13 indicate the sequences of the wild type and of mutant, respectively. SEQ ID NOS : 12 and 14 indicate amino acid sequences that these sequences can code for. Comparing the base sequences thus determined, it was confirmed that in the mutant serA, the 1087th base, G, was mutated into A and as a result, the 325th amino acid, glutamic acid, was changed to lysine.

(Example 6) Introduction of Plasmid Containing 3-PGDH Gene into Brevibacterium flavum

[0078] Plasmids harboring wild type serA or mutant serA were each introduced into Brevibacterium flavum AJ13377. The plasmids were introduced by the electric pulse method (Sugimoto et al., Japanese Patent Application Laid-open No. 2-207791). Transformed cells were selected in a complete medium containing 25 μ g/ml of kanamycin.

(Example 7) Production of L-serine by Transformed Cells

[0079] Transformed cells each having introduced therein plasmids harboring gene fragments containing wild serA or mutant serA in their full-length were incubated in a 500 ml shaking flask according to Example 3, and L-serine produced was determined. As a control, the AJ13377 strain as a host was incubated similarly.

[0080] In the transformed cell having introduced therein the wild type serA was observed no influence on its L-serine productivity whereas in the transformed cell having introduced therein the mutant serA was confirmed an increase in L-serine productivity (Table 3).

Table 3

Strain	Amplified Gene	Amount of L-serine that accumulated (g/L).
AJ13377	-	5.0
	serA	5.0

Table 3 (continued)

Strain	Amplified Gene	Amount of L-serine that accumulated (g/L)
	serA*	12.0

serA*: Mutant serA gene

[0081] Brevibacterium flavum AJ13377 has been deposited since October 15, 1997 in National Institute of Bioscience and Human Technology of Agency of Industrial Science and Technology of Ministry of International Trade and Industry (zip code: 305-8566, 1-3 Higashi 1-Chome, Tsukuba-shi, Ibaraki-ken, Japan), as accession number of FERM P-16471, and transferred from the original deposition to international deposition based on Budapest Treaty on November 20, 1998, and has been deposited as accession number of FERM BP-6576.

[0082] Further, the plasmid containing the mutant serA was harbored in Brevibacterium flavum ATCC 14067. The plasmid-harboring strain has been awarded Brevibacterium flavum AJ13378 and deposited since October 15, 1997 in National Institute of Bioscience and Human Technology of Agency of Industrial Science and Technology of Ministry of International Trade and Industry (zip code: 305-8566, 1-3 Higashi 1-Chome, Tsukuba-shi, Ibaraki-ken, Japan), as accession number of FERM P-16472, and transferred from the original deposition to international deposition based on Budapest Treaty on November 20, 1998, and has been deposited as accession number of FERM BP-6577.

Annex to the description

[0083]

SEQUENCE LISTING

<110> Ajinomoto Co., Inc
 <120> METHOD OF PRODUCING L-SERINE BY FERMENTATION
 <130> OP812
 <150> JP 10-3751
 <151> 1998-01-12
 <150> JP 10-353513
 <151> 1998-12-11
 <160> 14
 <170> PatentIn Ver. 2.0

 <210> 1
 <211> 1432
 <212> DNA
 <213> Corynebacterium glutamicum
 <220>
 <221> CDS
 <222> (398)..(1432)
 <400> 1
 ggatccggac acacgtgaca aaattgtaga aaattggatg attttgtcac gcctgtctgg 60
 tttagctctg gtteggggacg ggcgtggaat ggaggtagcg caccgagacc ttgacccgcy 120
 gcccgacaag ccaaaagtc ccaaaacaaa cccacctgc cygagacgtg aataaaattc 180
 gcagtcatt ccacagcgt aaacgcagct ttttgcattg tgagacacct ttgggggtaa 240
 atctcacagc atgaattctt ggggttagatg actttctggg tgggggaggg tttagaatgt 300
 ttctagtcgc acgccaaaac ccggcgtgga cacgtctgca gccgacgcgg tcgtgcctgt 360
 tgtaggcgga cattctagtg ttttccagga gtaactt gtg agc cag aat ggc cgt 415
 Val Ser Gln Asn Gly Arg
 1 5
 ccg gta gtc ctg atc gcc gat aag ctt gcg cag tcc act gtt gac gcg 463
 Pro Val Val Leu Ile Ala Asp Lys Leu Ala Gln Ser Thr Val Asp Ala
 10 15 20
 ctt gga gat gca gta gaa gtc cgt tgg gtt gac gga cct aac cgc cca 511
 Leu Gly Asp Ala Val Glu Val Arg Trp Val Asp Gly Pro Asn Arg Pro
 25 30 35
 gaa ctg ctt gat gca gtt aag gaa gcg gac gca ctg ctg gtg cgt tct 559
 Glu Leu Leu Asp Ala Val Lys Glu Ala Asp Ala Leu Leu Val Arg Ser
 40 45 50
 gct acc act gtc gat gct gaa gtc atc gcc gct gcc cct aac ttg aag 607
 Ala Thr Thr Val Asp Ala Glu Val Ile Ala Ala Pro Asn Leu Lys
 55 60 65 70

EP 0 943 687 A2

atc gtc ggt cgt gcc ggc gtg ggc ttg gac aac gtt gac atc cct gct 655
 Ile Val Gly Arg Ala Gly Val Gly Leu Asp Asn Val Asp Ile Pro Ala
 75 80 85
 gcc act gaa gct ggc gtc atg gtt gct aac gca ccg acc tct aac att 703
 Ala Thr Glu Ala Gly Val Met Val Ala Asn Ala Pro Thr Ser Asn Ile
 90 95 100
 cac tct gct tgt gag cac gca att tct ttg ctg ctg tct act gct cgc 751
 His Ser Ala Cys Glu His Ala Ile Ser Leu Leu Leu Ser Thr Ala Arg
 105 110 115
 cag atc cct gct gct gat gcg acg ctg cgt gag ggc gag tgg aag cgg 799
 Gln Ile Pro Ala Ala Asp Ala Thr Leu Arg Glu Gly Glu Trp Lys Arg
 120 125 130
 tct tct ttc aac ggt gtg gaa att ttc gga aaa act gtc ggt atc gtc 847
 Ser Ser Phe Asn Gly Val Glu Ile Phe Gly Lys Thr Val Gly Ile Val
 135 140 145 150
 ggt ttt ggc cac att ggt cag ttg ttt gct cag cgt ctt gct gcg ttt 895
 Gly Phe Gly His Ile Gly Gln Leu Phe Ala Gln Arg Leu Ala Ala Phe
 155 160 165
 gag acc acc att gtt gct tac gat cct tac gcc aac cct gct cgt gca 943
 Glu Thr Thr Ile Val Ala Tyr Asp Pro Tyr Ala Asn Pro Ala Arg Ala
 170 175 180
 gct cag ctg aac gtt gag ttg gtt gag ttg gat gag ctg atg agc cgt 991
 Ala Gln Leu Asn Val Glu Leu Val Glu Leu Asp Glu Leu Met Ser Arg
 185 190 195
 tct gac ttt gtc acc att cac ctt cct aag acc aag gaa act gct ggc 1039
 Ser Asp Phe Val Thr Ile His Leu Pro Lys Thr Lys Glu Thr Ala Gly
 200 205 210
 atg ttt gat gcg cag ctc ctt gct aag tcc aag aag ggc cag atc atc 1087
 Met Phe Asp Ala Gln Leu Leu Ala Lys Ser Lys Lys Gly Gln Ile Ile
 215 220 225 230
 atc aac gct gct cgt ggt ggc ctt gtt gat gag cag gct ttg gct gat 1135
 Ile Asn Ala Ala Arg Gly Gly Leu Val Asp Glu Gln Ala Leu Ala Asp
 235 240 245
 gcg att gag tcc ggt cac att cgt ggc gct ggt ttc gat gtg tac tcc 1183
 Ala Ile Glu Ser Gly His Ile Arg Gly Ala Gly Phe Asp Val Tyr Ser
 250 255 260
 acc gag cct tgc act gat tct cct ttg ttc aag ttg cct cag gtt gtt 1231
 Thr Glu Pro Cys Thr Asp Ser Pro Leu Phe Lys Leu Pro Gln Val Val
 265 270 275
 gtg act cct cac ttg ggt gct tct act gaa gag gct cag gat cgt gcg 1279
 Val Thr Pro His Leu Gly Ala Ser Thr Glu Glu Ala Gln Asp Arg Ala
 280 285 290

55

EP 0 943 687 A2

ggt act gac gtt gct gat tct gtg ctc aag gcg ctg gct ggc gag ttc 1327
 Gly Thr Asp Val Ala Asp Ser Val Leu Lys Ala Leu Ala Gly Glu Phe
 295 300 305 310
 gtg gcg gat gct gtg aac gtt tcc ggt ggt cgc gtg ggc gaa gag gtt 1375
 Val Ala Asp Ala Val Asn Val Ser Gly Gly Arg Val Gly Glu Glu Val
 315 320 325
 gct gtg tgg atg gat ctg gct cgc aag ctt ggt ctt ctt gct ggc aag 1423
 Ala Val Trp Met Asp Leu Ala Arg Lys Leu Gly Leu Leu Ala Gly Lys
 330 335 340
 ctt gtc gac 1432
 Leu Val Asp
 345

<210> 2

<211> 345

<212> PRT

<213> Corynebacterium glutamicum

<400> 2

Val Ser Gln Asn Gly Arg Pro Val Val Leu Ile Ala Asp Lys Leu Ala
 1 5 10 15
 Gln Ser Thr Val Asp Ala Leu Gly Asp Ala Val Glu Val Arg Trp Val
 20 25 30
 Asp Gly Pro Asn Arg Pro Glu Leu Leu Asp Ala Val Lys Glu Ala Asp
 35 40 45
 Ala Leu Leu Val Arg Ser Ala Thr Thr Val Asp Ala Glu Val Ile Ala
 50 55 60
 Ala Ala Pro Asn Leu Lys Ile Val Gly Arg Ala Gly Val Gly Leu Asp
 65 70 75 80
 Asn Val Asp Ile Pro Ala Ala Thr Glu Ala Gly Val Met Val Ala Asn
 85 90 95
 Ala Pro Thr Ser Asn Ile His Ser Ala Cys Glu His Ala Ile Ser Leu
 100 105 110
 Leu Leu Ser Thr Ala Arg Gln Ile Pro Ala Ala Asp Ala Thr Leu Arg
 115 120 125
 Glu Gly Glu Trp Lys Arg Ser Ser Phe Asn Gly Val Glu Ile Phe Gly
 130 135 140
 Lys Thr Val Gly Ile Val Gly Phe Gly His Ile Gly Gln Leu Phe Ala
 145 150 155 160
 Gln Arg Leu Ala Ala Phe Glu Thr Thr Ile Val Ala Tyr Asp Pro Tyr
 165 170 175
 Ala Asn Pro Ala Arg Ala Ala Gln Leu Asn Val Glu Leu Val Glu Leu
 180 185 190

Asp Glu Leu Met Ser Arg Ser Asp Phe Val Thr Ile His Leu Pro Lys
 195 200 205
 5 Thr Lys Glu Thr Ala Gly Met Phe Asp Ala Gln Leu Leu Ala Lys Ser
 210 215 220
 Lys Lys Gly Gln Ile Ile Ile Asn Ala Ala Arg Gly Gly Leu Val Asp
 225 230 235 240
 10 Glu Gln Ala Leu Ala Asp Ala Ile Glu Ser Gly His Ile Arg Gly Ala
 245 250 255
 Gly Phe Asp Val Tyr Ser Thr Glu Pro Cys Thr Asp Ser Pro Leu Phe
 260 265 270
 15 Lys Leu Pro Gln Val Val Val Thr Pro His Leu Gly Ala Ser Thr Glu
 275 280 285
 Glu Ala Gln Asp Arg Ala Gly Thr Asp Val Ala Asp Ser Val Leu Lys
 290 295 300
 20 Ala Leu Ala Gly Glu Phe Val Ala Asp Ala Val Asn Val Ser Gly Gly
 305 310 315 320
 Arg Val Gly Glu Glu Val Ala Val Trp Met Asp Leu Ala Arg Lys Leu
 325 330 335
 25 Gly Leu Leu Ala Gly Lys Leu Val Asp
 340 345

<210> 3

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Primer

<400> 3

ggacacacgt gacaaaattg tag

23

<210> 4

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Primer

<400> 4

gccagcaaga agaccaagct tgc

23

<210> 5

<211> 35

<212> DNA

5 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence:Primer
 <400> 5
 gtacatattg tcgtagaac gcgtaatacg actca 35

10 <210> 6
 <211> 23
 <212> DNA
 <213> Artificial Sequence
 <220>
 15 <223> Description of Artificial Sequence:Primer
 <400> 6
 tcatcaacgc tgctcgtggt ggc 23

20 <210> 7
 <211> 35
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <223> Description of Artificial Sequence:Primer
 <400> 7
 30 cgtagaacg cgtaatacga ctactatag ggaga 35

<210> 8
 <211> 23
 35 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence:Primer
 <400> 8
 40 gacgttgctg attctgtgct caa 23

45 <210> 9
 <211> 23
 <212> DNA
 <213> Artificial Sequence
 <220>
 50 <223> Description of Artificial Sequence:Primer
 <400> 9
 gggaggggttt agaattgttc tag 23

55

EP 0 943 687 A2

<210> 10

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Primer

<400> 10

ggttcaagca aatggatctc taa

23

<210> 11

<211> 1730

<212> DNA

<213> Brevibacterium flavum

<220>

<221> CDS

<222> (115)..(1704)

<400> 11

gggagggttt agaatgtttc tagtcgcacg ccaaaacccg gcgtggacac gtctgcagcc 60

gacgcggtcg tgctgttgt aggcggacat tctagtgtt tccaggagta actt gtg 117

Val

1

agc cag aat ggc cgt ccg gta gtc ctc atc gcc gat aag ctt gcg cag 165

Ser Gln Asn Gly Arg Pro Val Val Leu Ile Ala Asp Lys Leu Ala Gln

5

10

15

tcc act gtt gac gcg ctt gga gat gca gta gaa gtc cgt tgg gtt gac 213

Ser Thr Val Asp Ala Leu Gly Asp Ala Val Glu Val Arg Trp Val Asp

20

25

30

gga cct aac cgc cca gaa ctg ctt gat aca gtt aag gaa gcg gac gca 261

Gly Pro Asn Arg Pro Glu Leu Leu Asp Thr Val Lys Glu Ala Asp Ala

35

40

45

ctg ctc gtg cgt tct gct acc act gtc gat gct gaa gtc atc gcc gct 309

Leu Leu Val Arg Ser Ala Thr Thr Val Asp Ala Glu Val Ile Ala Ala

50

55

60

65

gcc cct aac ttg aag atc gtc ggt cgt gcc gcc gtg gcc ttg gac aac 357

Ala Pro Asn Leu Lys Ile Val Gly Arg Ala Gly Val Gly Leu Asp Asn

70

75

80

gtt gac atc cct gct gcc act gaa gct gcc gtc atg gtt gct aac gca 405

Val Asp Ile Pro Ala Ala Thr Glu Ala Gly Val Met Val Ala Asn Ala

85

90

95

ccg acc tct aac att cac tct gct tgt gag cac gca att tct ttg ctg 453

Pro Thr Ser Asn Ile His Ser Ala Cys Glu His Ala Ile Ser Leu Leu

100

105

110

EP 0 943 687 A2

	ctg tct act gct cgc cag atc cct gct gct gat gcg acg ctg cgt gag	501
	Leu Ser Thr Ala Arg Gln Ile Pro Ala Ala Asp Ala Thr Leu Arg Glu	
5	115 120 125	
	ggc gag tgg aag cgg tct tct ttc aac ggt gtg gaa att ttc gga aaa	549
	Gly Glu Trp Lys Arg Ser Ser Phe Asn Gly Val Glu Ile Phe Gly Lys	
	130 135 140 145	
10	act gtc ggt atc gtc ggt ttt ggc cac att ggt cag ttg ttt gct cag	597
	Thr Val Gly Ile Val Gly Phe Gly His Ile Gly Gln Leu Phe Ala Gln	
	150 155 160	
	cgt ctt gct gcg ttt gag acc acc att gtt gct tac gat cct tac gct	645
15	Arg Leu Ala Ala Phe Glu Thr Thr Ile Val Ala Tyr Asp Pro Tyr Ala	
	165 170 175	
	aac cct gct cgt gcg gct cag ctg aac gtt gag ttg gtt gag ttg gat	693
	Asn Pro Ala Arg Ala Ala Gln Leu Asn Val Glu Leu Val Glu Leu Asp	
	180 185 190	
20	gag ctg atg agc cgt tct gac ttt gtc acc att cac ctt cct aag acc	741
	Glu Leu Met Ser Arg Ser Asp Phe Val Thr Ile His Leu Pro Lys Thr	
	195 200 205	
	aag gaa act gct ggc atg ttt gat gcg cag ctc ctt gct aag tcc aag	789
25	Lys Glu Thr Ala Gly Met Phe Asp Ala Gln Leu Leu Ala Lys Ser Lys	
	210 215 220 225	
	aag ggc cag atc atc atc aac gct gct cgt ggt ggc ctt gtt gat gaa	837
	Lys Gly Gln Ile Ile Ile Asn Ala Ala Arg Gly Gly Leu Val Asp Glu	
30	230 235 240	
	cag gct ttg gct gat gcg att gag tcc ggt cac att cgt ggc gct ggt	885
	Gln Ala Leu Ala Asp Ala Ile Glu Ser Gly His Ile Arg Gly Ala Gly	
	245 250 255	
35	ttc gat gtg tac tcc acc gag cct tgc act gat tct cct ttg ttc aag	933
	Phe Asp Val Tyr Ser Thr Glu Pro Cys Thr Asp Ser Pro Leu Phe Lys	
	260 265 270	
	ttg cct cag gtt gtt gtg act cct cac ttg ggt gct tct act gaa gag	981
40	Leu Pro Gln Val Val Val Thr Pro His Leu Gly Ala Ser Thr Glu Glu	
	275 280 285	
	gct cag gat cgt gcg ggt act gac gtt gct gat tct gtg ctc aag gcg	1029
	Ala Gln Asp Arg Ala Gly Thr Asp Val Ala Asp Ser Val Leu Lys Ala	
45	290 295 300 305	
	ctg gct ggc gag ttc gtg gcg gat gct gtg aac gtt tcc ggt ggt cgc	1077
	Leu Ala Gly Glu Phe Val Ala Asp Ala Val Asn Val Ser Gly Gly Arg	
	310 315 320	
50	gtg ggc gaa gag gtt gct gtg tgg atg gat ctg gct cgc aag ctt ggt	1125
	Val Gly Glu Glu Val Ala Val Trp Met Asp Leu Ala Arg Lys Leu Gly	
	325 330 335	

EP 0 943 687 A2

5 ctt ctt gct ggc aag ctt gtc gac gcc gcc cca gtc tcc att gag gtt 1173
 Leu Leu Ala Gly Lys Leu Val Asp Ala Ala Pro Val Ser Ile Glu Val
 340 345 350
 gag gct cga ggc gag ctt tct tcc gag cag gtc gat gca ctt ggt ttg 1221
 Glu Ala Arg Gly Glu Leu Ser Ser Glu Gln Val Asp Ala Leu Gly Leu
 355 360 365
 10 tcc gct gtt cgt ggt ttg ttc tcc gga att atc gaa gag tcc gtt act 1269
 Ser Ala Val Arg Gly Leu Phe Ser Gly Ile Ile Glu Glu Ser Val Thr
 370 375 380 385
 ttc gtc aac gct cct cgc att gct gaa gag cgt ggc ctg gac atc tcc 1317
 Phe Val Asn Ala Pro Arg Ile Ala Glu Glu Arg Gly Leu Asp Ile Ser
 390 395 400
 gtg aag acc aac tct gag tct gtt act cac cgt tcc gtc ctg cag gtc 1365
 Val Lys Thr Asn Ser Glu Ser Val Thr His Arg Ser Val Leu Gln Val
 405 410 415
 20 aag gtc att act ggc agc ggc gcg agc gca act gtt gtt ggt gcc ctg 1413
 Lys Val Ile Thr Gly Ser Gly Ala Ser Ala Thr Val Val Gly Ala Leu
 420 425 430
 act ggt ctt gag cgc gtt gag aag atc acc cgc atc aat ggc cgt ggc 1461
 Thr Gly Leu Glu Arg Val Glu Lys Ile Thr Arg Ile Asn Gly Arg Gly
 435 440 445
 ctg gat ctg cgc gca gag ggt ctg aac ctc ttc ctg cag tac act gac 1509
 Leu Asp Leu Arg Ala Glu Gly Leu Asn Leu Phe Leu Gln Tyr Thr Asp
 450 455 460 465
 gct cct ggt gca ctg ggt acc gtt ggt acc aag ctg ggt gct gct ggc 1557
 Ala Pro Gly Ala Leu Gly Thr Val Gly Thr Lys Leu Gly Ala Ala Gly
 470 475 480
 35 atc aac atc gag gct gct gcg ttg act cag gct gag aag ggt gac ggc 1605
 Ile Asn Ile Glu Ala Ala Ala Leu Thr Gln Ala Glu Lys Gly Asp Gly
 485 490 495
 gct gtc ctg atc ctg cgt gtt gag tcc gct gtc tcc gaa gag ctg gaa 1653
 Ala Val Leu Ile Leu Arg Val Glu Ser Ala Val Ser Glu Glu Leu Glu
 500 505 510
 gct gaa atc aac gct gag ttg ggt gct act tcc ttc cag gtt gat ctt 1701
 Ala Glu Ile Asn Ala Glu Leu Gly Ala Thr Ser Phe Gln Val Asp Leu
 515 520 525
 45 gac taattagaga tccattttct agaacc 1730
 Asp
 530
 50 <210> 12
 <211> 530
 55

<212> PRT

<213> Brevibacterium flavum

<400> 12

5 Val Ser Gln Asn Gly Arg Pro Val Val Leu Ile Ala Asp Lys Leu Ala
 1 5 10 15
 Gln Ser Thr Val Asp Ala Leu Gly Asp Ala Val Glu Val Arg Trp Val
 20 25 30
 10 Asp Gly Pro Asn Arg Pro Glu Leu Leu Asp Thr Val Lys Glu Ala Asp
 35 40 45
 Ala Leu Leu Val Arg Ser Ala Thr Thr Val Asp Ala Glu Val Ile Ala
 50 55 60
 15 Ala Ala Pro Asn Leu Lys Ile Val Gly Arg Ala Gly Val Gly Leu Asp
 65 70 75 80
 Asn Val Asp Ile Pro Ala Ala Thr Glu Ala Gly Val Met Val Ala Asn
 85 90 95
 20 Ala Pro Thr Ser Asn Ile His Ser Ala Cys Glu His Ala Ile Ser Leu
 100 105 110
 Leu Leu Ser Thr Ala Arg Gln Ile Pro Ala Ala Asp Ala Thr Leu Arg
 115 120 125
 25 Glu Gly Glu Trp Lys Arg Ser Ser Phe Asn Gly Val Glu Ile Phe Gly
 130 135 140
 Lys Thr Val Gly Ile Val Gly Phe Gly His Ile Gly Gln Leu Phe Ala
 145 150 155 160
 30 Gln Arg Leu Ala Ala Phe Glu Thr Thr Ile Val Ala Tyr Asp Pro Tyr
 165 170 175
 Ala Asn Pro Ala Arg Ala Ala Gln Leu Asn Val Glu Leu Val Glu Leu
 180 185 190
 35 Asp Glu Leu Met Ser Arg Ser Asp Phe Val Thr Ile His Leu Pro Lys
 195 200 205
 Thr Lys Glu Thr Ala Gly Met Phe Asp Ala Gln Leu Leu Ala Lys Ser
 210 215 220
 40 Lys Lys Gly Gln Ile Ile Ile Asn Ala Ala Arg Gly Gly Leu Val Asp
 225 230 235 240
 Glu Gln Ala Leu Ala Asp Ala Ile Glu Ser Gly His Ile Arg Gly Ala
 245 250 255
 45 Gly Phe Asp Val Tyr Ser Thr Glu Pro Cys Thr Asp Ser Pro Leu Phe
 260 265 270
 Lys Leu Pro Gln Val Val Val Thr Pro His Leu Gly Ala Ser Thr Glu
 275 280 285
 50 Glu Ala Gln Asp Arg Ala Gly Thr Asp Val Ala Asp Ser Val Leu Lys
 290 295 300
 55 Ala Leu Ala Gly Glu Phe Val Ala Asp Ala Val Asn Val Ser Gly Gly

EP 0 943 687 A2

305 310 315 320
 Arg Val Gly Glu Glu Val Ala Val Trp Met Asp Leu Ala Arg Lys Leu
 5 325 330 335
 Gly Leu Leu Ala Gly Lys Leu Val Asp Ala Ala Pro Val Ser Ile Glu
 340 345 350
 Val Glu Ala Arg Gly Glu Leu Ser Ser Glu Gln Val Asp Ala Leu Gly
 10 355 360 365
 Leu Ser Ala Val Arg Gly Leu Phe Ser Gly Ile Ile Glu Glu Ser Val
 370 375 380
 Thr Phe Val Asn Ala Pro Arg Ile Ala Glu Glu Arg Gly Leu Asp Ile
 15 385 390 395 400
 Ser Val Lys Thr Asn Ser Glu Ser Val Thr His Arg Ser Val Leu Gln
 405 410 415
 Val Lys Val Ile Thr Gly Ser Gly Ala Ser Ala Thr Val Val Gly Ala
 20 420 425 430
 Leu Thr Gly Leu Glu Arg Val Glu Lys Ile Thr Arg Ile Asn Gly Arg
 435 440 445
 Gly Leu Asp Leu Arg Ala Glu Gly Leu Asn Leu Phe Leu Gln Tyr Thr
 25 450 455 460
 Asp Ala Pro Gly Ala Leu Gly Thr Val Gly Thr Lys Leu Gly Ala Ala
 465 470 475 480
 Gly Ile Asn Ile Glu Ala Ala Ala Leu Thr Gln Ala Glu Lys Gly Asp
 30 485 490 495
 Gly Ala Val Leu Ile Leu Arg Val Glu Ser Ala Val Ser Glu Glu Leu
 500 505 510
 Glu Ala Glu Ile Asn Ala Glu Leu Gly Ala Thr Ser Phe Gln Val Asp
 35 515 520 525
 Leu Asp
 530
 40 <210> 13
 <211> 173G
 <212> DNA
 <213> Brevibacterium flavum
 45 <220>
 <221> CDS
 <222> (115)..(1704)
 <400> 13
 50 gggaggggttt agaatgtttc tagtcgcacg ccaaaaccg gcgtggacac gtctgcagcc 60
 gacgcggtcg tgcctgttgt aggcggacat tcctagtttt tccaggagta actt gtg 117
 Val
 55 1

EP 0 943 687 A2

agc cag aat ggc cgt ccg gta gtc etc atc gcc gat aag ctt gcg cag 165
 Ser Gln Asn Gly Arg Pro Val Val Leu Ile Ala Asp Lys Leu Ala Gln
 5 5 10 15
 tcc act gtt gac gcg ctt gga gat gca gta gaa gtc cgt tgg gtt gac 213
 Ser Thr Val Asp Ala Leu Gly Asp Ala Val Glu Val Arg Trp Val Asp
 20 25 30
 gga cct aac cgc cca gaa ctg ctt gat aca gtt aag gaa gcg gac gca 261
 Gly Pro Asn Arg Pro Glu Leu Leu Asp Thr Val Lys Glu Ala Asp Ala
 35 40 45
 ctg ctc gtg cgt tct gct acc act gtc gat gct gaa gtc atc gcc gct 309
 Leu Leu Val Arg Ser Ala Thr Thr Val Asp Ala Glu Val Ile Ala Ala
 50 55 60 65
 gcc cct aac ttg aag atc gtc ggt cgt gcc ggc gtg ggc ttg gac aac 357
 Ala Pro Asn Leu Lys Ile Val Gly Arg Ala Gly Val Gly Leu Asp Asn
 70 75 80
 gtt gac atc cct gct gcc act gaa gct ggc gtc atg gtt gct aac gca 405
 Val Asp Ile Pro Ala Ala Thr Glu Ala Gly Val Met Val Ala Asn Ala
 85 90 95
 ccg acc tct aac att cac tct gct tgt gag cac gca att tct ttg ctg 453
 Pro Thr Ser Asn Ile His Ser Ala Cys Glu His Ala Ile Ser Leu Leu
 100 105 110
 ctg tct act gct cgc cag atc cct gct gct gat gcg acg ctg cgt gag 501
 Leu Ser Thr Ala Arg Gln Ile Pro Ala Ala Asp Ala Thr Leu Arg Glu
 115 120 125
 ggc gag tgg aag cgg tct tct ttc aac ggt gtg gaa att ttc gga aaa 549
 Gly Glu Trp Lys Arg Ser Ser Phe Asn Gly Val Glu Ile Phe Gly Lys
 130 135 140 145
 act gtc ggt atc gtc ggt ttt ggc cac att ggt cag ttg ttt gct cag 597
 Thr Val Gly Ile Val Gly Phe Gly His Ile Gly Gln Leu Phe Ala Gln
 150 155 160
 cgt ctt gct gcg ttt gag acc acc att gtt gct tac gat cct tac gct 645
 Arg Leu Ala Ala Phe Glu Thr Thr Ile Val Ala Tyr Asp Pro Tyr Ala
 165 170 175
 aac cct gct cgt gcg gct cag ctg aac gtt gag ttg gtt gag ttg gat 693
 Asn Pro Ala Arg Ala Ala Gln Leu Asn Val Glu Leu Val Glu Leu Asp
 180 185 190
 gag ctg atg agc cgt tct gac ttt gtc acc att cac ctt cct aag acc 741
 Glu Leu Met Ser Arg Ser Asp Phe Val Thr Ile His Leu Pro Lys Thr
 195 200 205
 aag gaa act gct ggc atg ttt gat gcg cag etc ctt gct aag tcc aag 789
 Lys Glu Thr Ala Gly Met Phe Asp Ala Gln Leu Leu Ala Lys Ser Lys
 210 215 220 225

EP 0 943 687 A2

aag ggc cag atc atc atc aac gct gct cgt ggt ggc ctt gtt gat gaa 837
 Lys Gly Gln Ile Ile Ile Asn Ala Ala Arg Gly Gly Leu Val Asp Glu
 230 235 240
 5 cag gct ttg gct gat gcg att gag tcc ggt cac att cgt ggc gct ggt 885
 Gln Ala Leu Ala Asp Ala Ile Glu Ser Gly His Ile Arg Gly Ala Gly
 245 250 255
 10 ttc gat gtg tac tcc acc gag cct tgc act gat tct cct ttg ttc aag 933
 Phe Asp Val Tyr Ser Thr Glu Pro Cys Thr Asp Ser Pro Leu Phe Lys
 260 265 270
 ttg cct cag gtt gtt gtg act cct cac ttg ggt gct tct act gaa gag 981
 Leu Pro Gln Val Val Val Thr Pro His Leu Gly Ala Ser Thr Glu Glu
 15 275 280 285
 gct cag gat cgt gcg ggt act gac gtt gct gat tct gtg ctc aag gcg 1029
 Ala Gln Asp Arg Ala Gly Thr Asp Val Ala Asp Ser Val Leu Lys Ala
 290 295 300 305
 20 ctg gct ggc gag ttc gtg gcg gat gct gtg aac gtt tcc ggt ggt cgc 1077
 Leu Ala Gly Glu Phe Val Ala Asp Ala Val Asn Val Ser Gly Gly Arg
 310 315 320
 gtg ggc gaa aag gtt gct gtg tgg atg gat ctg gct cgc aag ctt ggt 1125
 25 Val Gly Glu Lys Val Ala Val Trp Met Asp Leu Ala Arg Lys Leu Gly
 325 330 335
 ctt ctt gct ggc aag ctt gtc gac gcc gcc cca gtc tcc att gag gtt 1173
 Leu Leu Ala Gly Lys Leu Val Asp Ala Ala Pro Val Ser Ile Glu Val
 340 345 350
 30 gag gct cga ggc gag ctt tct tcc gag cag gtc gat gca ctt ggt ttg 1221
 Glu Ala Arg Gly Glu Leu Ser Ser Glu Gln Val Asp Ala Leu Gly Leu
 355 360 365
 tcc gct gtt cgt ggt ttg ttc tcc gga att atc gaa gag tcc gtt act 1269
 35 Ser Ala Val Arg Gly Leu Phe Ser Gly Ile Ile Glu Glu Ser Val Thr
 370 375 380 385
 ttc gtc aac gct cct cgc att gct gaa gag cgt ggc ctg gac atc tcc 1317
 Phe Val Asn Ala Pro Arg Ile Ala Glu Glu Arg Gly Leu Asp Ile Ser
 390 395 400
 40 gtg aag acc aac tct gag tct gtt act cac cgt tcc gtc ctg cag gtc 1365
 Val Lys Thr Asn Ser Glu Ser Val Thr His Arg Ser Val Leu Gln Val
 405 410 415
 45 aag gtc att act ggc agc ggc gcg agc gca act gtt gtt ggt gcc ctg 1413
 Lys Val Ile Thr Gly Ser Gly Ala Ser Ala Thr Val Val Gly Ala Leu
 420 425 430
 act ggt ctt gag cgc gtt gag aag atc acc cgc atc aat ggc cgt ggc 1461
 50 Thr Gly Leu Glu Arg Val Glu Lys Ile Thr Arg Ile Asn Gly Arg Gly
 435 440 445

EP 0 943 687 A2

ctg gat ctg cgc gca gag ggt ctg aac ctc ttc ctg cag tac act gac 1509
 Leu Asp Leu Arg Ala Glu Gly Leu Asn Leu Phe Leu Gln Tyr Thr Asp
 450 455 460 465
 5 gct cct ggt gca ctg ggt acc gtt ggt acc aag ctg ggt gct gct ggc 1557
 Ala Pro Gly Ala Leu Gly Thr Val Gly Thr Lys Leu Gly Ala Ala Gly
 470 475 480
 10 atc aac atc gag gct gct gcg ttg act cag gct gag aag ggt gac ggc 1605
 Ile Asn Ile Glu Ala Ala Ala Leu Thr Gln Ala Glu Lys Gly Asp Gly
 485 490 495
 15 gct gtc ctg atc ctg cgt gtt gag tcc gct gtc tcc gaa gag ctg gaa 1653
 Ala Val Leu Ile Leu Arg Val Glu Ser Ala Val Ser Glu Glu Leu Glu
 500 505 510
 gct gaa atc aac gct gag ttg ggt gct act tcc ttc cag gtt gat ctt 1701
 Ala Glu Ile Asn Ala Glu Leu Gly Ala Thr Ser Phe Gln Val Asp Leu
 515 520 525
 20 gac taattagaga tccattttct agaacc 1730
 Asp
 530
 25 <210> 14
 <211> 530
 <212> PRT
 <213> Brevibacterium flavum
 30 <400> 14
 Val Ser Gln Asn Gly Arg Pro Val Val Leu Ile Ala Asp Lys Leu Ala
 1 5 10 15
 35 Gln Ser Thr Val Asp Ala Leu Gly Asp Ala Val Glu Val Arg Trp Val
 20 25 30
 Asp Gly Pro Asn Arg Pro Glu Leu Leu Asp Thr Val Lys Glu Ala Asp
 35 40 45
 40 Ala Leu Leu Val Arg Ser Ala Thr Thr Val Asp Ala Glu Val Ile Ala
 50 55 60
 Ala Ala Pro Asn Leu Lys Ile Val Gly Arg Ala Gly Val Gly Leu Asp
 65 70 75 80
 45 Asn Val Asp Ile Pro Ala Ala Thr Glu Ala Gly Val Met Val Ala Asn
 85 90 95
 Ala Pro Thr Ser Asn Ile His Ser Ala Cys Glu His Ala Ile Ser Leu
 100 105 110
 50 Leu Leu Ser Thr Ala Arg Gln Ile Pro Ala Ala Asp Ala Thr Leu Arg
 115 120 125
 Glu Gly Glu Trp Lys Arg Ser Ser Phe Asn Gly Val Glu Ile Phe Gly
 130 135 140
 55

Lys Thr Val Gly Ile Val Gly Phe Gly His Ile Gly Gln Leu Phe Ala
 145 150 155 160
 5 Gln Arg Leu Ala Ala Phe Glu Thr Thr Ile Val Ala Tyr Asp Pro Tyr
 165 170 175
 Ala Asn Pro Ala Arg Ala Ala Gln Leu Asn Val Glu Leu Val Glu Leu
 180 185 190
 10 Asp Glu Leu Met Ser Arg Ser Asp Phe Val Thr Ile His Leu Pro Lys
 195 200 205
 Thr Lys Glu Thr Ala Gly Met Phe Asp Ala Gln Leu Leu Ala Lys Ser
 210 215 220
 15 Lys Lys Gly Gln Ile Ile Ile Asn Ala Ala Arg Gly Gly Leu Val Asp
 225 230 235 240
 Glu Gln Ala Leu Ala Asp Ala Ile Glu Ser Gly His Ile Arg Gly Ala
 245 250 255
 20 Gly Phe Asp Val Tyr Ser Thr Glu Pro Cys Thr Asp Ser Pro Leu Phe
 260 265 270
 Lys Leu Pro Gln Val Val Val Thr Pro His Leu Gly Ala Ser Thr Glu
 275 280 285
 25 Glu Ala Gln Asp Arg Ala Gly Thr Asp Val Ala Asp Ser Val Leu Lys
 290 295 300
 Ala Leu Ala Gly Glu Phe Val Ala Asp Ala Val Asn Val Ser Gly Gly
 305 310 315 320
 30 Arg Val Gly Glu Lys Val Ala Val Trp Met Asp Leu Ala Arg Lys Leu
 325 330 335
 Gly Leu Leu Ala Gly Lys Leu Val Asp Ala Ala Pro Val Ser Ile Glu
 340 345 350
 35 Val Glu Ala Arg Gly Glu Leu Ser Ser Glu Gln Val Asp Ala Leu Gly
 355 360 365
 Leu Ser Ala Val Arg Gly Leu Phe Ser Gly Ile Ile Glu Glu Ser Val
 370 375 380
 40 Thr Phe Val Asn Ala Pro Arg Ile Ala Glu Glu Arg Gly Leu Asp Ile
 385 390 395 400
 Ser Val Lys Thr Asn Ser Glu Ser Val Thr His Arg Ser Val Leu Gln
 405 410 415
 45 Val Lys Val Ile Thr Gly Ser Gly Ala Ser Ala Thr Val Val Gly Ala
 420 425 430
 Leu Thr Gly Leu Glu Arg Val Glu Lys Ile Thr Arg Ile Asn Gly Arg
 435 440 445
 50 Gly Leu Asp Leu Arg Ala Glu Gly Leu Asn Leu Phe Leu Gln Tyr Thr
 450 455 460
 Asp Ala Pro Gly Ala Leu Gly Thr Val Gly Thr Lys Leu Gly Ala Ala
 465 470 475 480

Gly Ile Asn Ile Glu Ala Ala Ala Leu Thr Gln Ala Glu Lys Gly Asp
 485 490 495
 5 Gly Ala Val Leu Ile Leu Arg Val Glu Ser Ala Val Ser Glu Glu Leu
 500 505 510
 Glu Ala Glu Ile Asn Ala Glu Leu Gly Ala Thr Ser Phe Gln Val Asp
 515 520 525
 10 Leu Asp
 530

15 Claims

1. A coryneform bacterium having resistance to azaserine or β -(2-thienyl)-DL-alanine and having L-serine productivity.
- 20 2. D-3-phosphoglycerate dehydrogenase derived from a coryneform bacterium, in which feedback inhibition by L-serine is desensitized.
3. The D-3-phosphoglycerate dehydrogenase as claimed in claim 2, which is obtainable from a coryneform bacterium having resistance to azaserine or β -(2-thienyl)-DL-alanine and having L-serine productivity.
- 25 4. The D-3-phosphoglycerate dehydrogenase as claimed in claim 2, having an amino acid sequence depicted in SEQ ID NO : 12 in Sequence Listing or said sequence including substitution, addition or deletion of one or more amino acids, wherein an amino acid residue corresponding to the 325th glutamic acid residue of the amino acid sequence in the SEQ ID NO : 12 is replaced with an amino acid other than glutamic acid.
- 30 5. The D-3-phosphoglycerate dehydrogenase as claimed in claim 2, which has an amino acid sequence depicted in SEQ ID NO : 11 in Sequence Listing.
- 35 6. A DNA coding for the D-3-phosphoglycerate dehydrogenase as claimed in any one of claims 2 to 5.
7. The DNA as claimed in claim 6, wherein said DNA has a base sequence depicted in SEQ ID NO : 13 in Sequence Listing.
- 40 8. A coryneform bacterium which harbors a recombinant DNA containing the DNA as claimed in claim 6.
9. A method of producing L-serine, comprising the steps of cultivating the bacterium as claimed in claim 1 or 8 in a medium to allow accumulation of L-serine in the medium, and collecting the L-serine from the medium.




FIG. 1




FIG. 2

THIS PAGE BLANK (USPTO)